

**WELLISAIR**

**VS**

**CORONAVIRUS**

**COVID-19**

## Effectiveness of the Wellisair Model WADU-02 Disinfection Purifier Against SARS-CoV-2 Contaminated Surfaces

This report was prepared at MRIGlobal for the work performed under MRIGlobal Task No. 311732.01.001, “Effectiveness of the Wellisair Model WADU-02 Disinfection Purifier against SARS-CoV-2 Contaminated Surfaces.”

The experimental phase of this task was initiated by MRIGlobal on April 20, 2021 and ended on April 23, 2021.

### Objective:

The objective of this project was to determine if the Wellisair Model WADU-02 Disinfection Purifier device has the ability to decrease viral infectivity of SARS-CoV-2 *in vitro* after exposure. The device was tested on SARS-CoV-2 virus inoculated on stainless steel coupons.

### Study Design:

Stainless steel coupons were inoculated with 200  $\mu$ L virus stock ( $2.37E5$  TCID<sub>50</sub>/ml of Washington 1 USA SARS-CoV-2 isolate). Virus was evenly spread over the coupons and allowed to dry. Test coupons were transferred to the testing room and placed into an aerosol test system inside a Class III Biosafety and exposed to Wellisair Model WADU-02. After the exposure time, any remaining virus was resuspended with a cell scraper and 2 ml Dulbecco’s Modified Eagle Medium with supplement F-12 (DMEM/F12). Samples were diluted 1:10 down a 96 deep well plate in DMEM/F12. These dilutions were transferred to a plate of Vero E6 cells with media removed. After approximately 45 minutes, DMEM/F12 supplemented with 5% fetal bovine serum (FBS) was added to cells to feed them for the next three days. The inoculated plates were then read for cytopathic effects (CPE).

### Results

Plates were read 3 days after the initiation of the assay. Wellisair Model WADU-02 exposure resulted in a log reduction of 1.08 compared to controls. Thus, in two hours 91.75% of SARS-CoV-2 infectivity was reduced *in vitro*.


**Table 1. Results of viral exposure to Wellisair Model WADU-02**

Sample Name	Test Article	Description	TCID <sub>50</sub> /ml	Log 10 TCID <sub>50</sub> /ml	avg TCID <sub>50</sub> /ml	avg log <sub>10</sub> TCID <sub>50</sub> /ml	Log Reduction	% Reduction
T1	Wellisair WADU-02	Test	6.81E+03	3.83	8.78E+03	3.92	1.08	91.75%
T2			6.81E+03	3.83				
T3			1.47E+04	4.17				
T4			6.81E+03	3.83				
C1		Control	4.22E+04	4.63	1.24E+05	5.00		
C2			1.47E+05	5.17				
C3			6.81E+04	4.83				
C4			2.37E+05	5.38				

### Conclusions:

Based on these experiments, viral infectivity was reduced by 1.08 log (91.75%) after exposure to Wellisair Model WADU-02 for two hours when compared to control samples.


## Wellisair versus Coronavirus - COVID-19




To understand the effectiveness of Wellisair against the COVID-19 pandemic (initially 2019-nCoV) that emerged in 2019 in Wuhan, one must understand the morphological structure of this type of virus, as well as its classification.

### Morphological structure and classification

Coronaviruses are in the Coronavirinae subfamily in the Coronaviridae family, in the order Nidovirales. They are divided into 4 VOC subgenera: Alpha, Beta, Delta and Gamma. COVID-19 belongs to the SARS family, classified as Betacoronavirus of lineage "B", originating in bats.




Based on the genetic material, these viruses are included in group IV of the Baltimore classification, as the viral particle contains only one strand of positive polarity linear single-stranded RNA.



The diameter of the virus is around 60 to 200nm, they present a nucleocapsid with helical symmetry and a lipid sheath derived from the membrane of the previously infected host cell. From these pods arise the characteristic projections in the form of a solar corona.

Given the short time since the appearance of this strain, direct tests of Wellisair technology against the virus have not been possible.

But it has been possible to test a virus with a morphological structure similar to the coronavirus, such as the Respiratory Syncytial Virus (RSV). RSV is classified within the Pneumovirus, specifically in the Paramyxoviridae family, and although they are not within the same family, this virus shares great similarities with COVID-19, such as the lipid membrane and glycoprotein projections.



The oxidation reactions (lipid oxidation and protein oxidation) that occur in RSV are the same as they should have in COVID-19, so the elimination efficiency should be very similar.

Based on the results obtained with Wellisair in viruses with similar structures (RSV) to Coronavirus COVID-19, we can expect that the effectiveness of our new technology will have an expected average elimination result of 92 to 99% depending on the relative humidity conditions.

Pathogen	Test result			Laboratory	Report No.
	Medium	Exhibition time	Effectiveness %		
Synchytial VRS	Wet	2 hours	99	Laboratory of viruses contaminating water and food University of Barcelona	20191212_3
	Dry	2 hours	92		20191212_4

These conclusions are included in a study carried out by researchers from the Center for Research on Food Safety and Control (CRESCA) of the Polytechnic University of Catalonia (UPC) and the e-SHealth published in the IJORS journal "International Journal of Engineering Research & Science".

### Bibliographic Report University of Barcelona

The microbiology, virology and biotechnology section of the water and food contaminating virus laboratory of the University of Barcelona has produced a bibliographic report on disinfection and the Coronavirus COVID-19.

Coronaviruses that are most similar to SARS-CoV are easily inactivated with commonly used disinfectants, and especially are más susceptibles a aquellos con base alcohólica more susceptible to alcohol-based ones (such as OH hydroxyl radical technology).

The Coronavirus has a morphology very similar to the RSV respiratory syncytial virus tested by the University of Barcelona with a reduction power of 92 to 99% (dry/wet) in 2 hours.

Although ozone is an especially efficient disinfectant for RNA viruses, high concentrations are harmful to living things and therefore not suitable for human-inhabited environments.

On the other hand, with the hydroxyl radicals OH generated by Wellisair, according to the results obtained in our UB laboratory, a similar activation of the Coronavirus is expected without the disadvantages of ozone toxicity.

**WELLISAIR**  
**Laboratory & University Test Results**  
**Pathogen – VIRUS**

Pathogen	Test result				
VIRUS	Medium	Exposure	Effectiveness %	Laboratory / University	Report N°
Influenzavirus	Wet	30 min	86	Laboratory of Viruses Contaminants of water and food University of Barcelona	20200228
	Dry	30 min	38		
Sincitial VRS	Wet	2 h	99	Laboratory of Viruses Contaminants of water and food University of Barcelona	20191212_3
	Dry	2 h	92		20191212_4
Rotavirus	Dry	2h	99	Laboratory of Viruses Contaminants of water and food University of Barcelona	20191212_2
Coxiackievirus (H2O2)	Wet	4 h	99.9	Laboratory of Viruses Contaminants of water and food University of Barcelona	20180711
	Dry	4 h	99.4		

**WELLISAIR**  
**Laboratory & University Test Results**  
**Pathogen – BACTERIA**

Pathogen	Test result				
VIRUS	Medium	Exposure	Effectiveness %	Laboratory / University	Report N°
Bacillus subtilis	Surface	1h	99.4	Kangwon National University (Korea)	EM20150002
	Air	20min	99.6	Kangwon National University (Korea)	EM20150002
Staphylococcus aureus	Surface	2h	52.3	Kangwon National University (Korea)	EM20150002
	Surface	2h	99.9	Korea Conformity Laboratories	027128
	Air	1h	99.9	Kangwon National University (Korea)	EM20150002
Staphylococcus aureus resistant MRSA	Surface	4h	99.9	Korea Conformity Laboratories	027129
	Air	4h	99.9	Korea Conformity Laboratories	027131
Pseudomonas aeruginosa	Surface	4h	99.9	Korea Conformity Laboratories	027128
Enterobacter species: Salmonella	Surface	4h	99.9	Korea Conformity Laboratories	027129
Enterobacter species: Klebsiella	Surface	4h	99.9	Korea Conformity Laboratories	027129
	Air	4h	99.9	Korea Conformity Laboratories	027130
Enterobacter species: Escherichia coli	Surface	1h	99.9	Kangwon National University (Korea)	EM20150002
	Surface	4h	99.9	Korea Conformity Laboratories	027128
	Air	20min	99.9	Kangwon National University (Korea)	EM20150002

**WELLISAIR**  
**Laboratory & University Test Results**  
**Air Quality: Volatile Organic Compounds (VOCs) & Atmospheric Particles**

Air Quality: Atmospheric pollutants 94		Test result		Laboratory / University	Nº Informe
		Exposure	Effectiveness %		
Volatile Organic Compounds	Total	30min	94.3	CRESCA (Polytechnic University of Catalonia)	20190930
	Formaldehyde	30min	93.4		
Atmospheric Particles	PM 2,5	30min	94.7	CRESCA (Polytechnic University of Catalonia)	20190930
	PM 1	30min	95.3		
	PM 10	30min	95.4		

**WELLISAIR**  
**Laboratory & University Test Results**  
**Air Quality: Deodorization Volatile Organic Compounds (VOCs)**

Air Quality: deodorization		Test result		Laboratory / University	Nº Informe
		Exposure	Effectiveness %		
Volatile Organic Compounds	Formaldehyde	4h	65.0	Korea Conformity Laboratories	026961
	Ammoniac	4h	80.0		
	Toluene	4h	99.5		

**WELLISAIR**  
**Cases Study Test Results**  
**Pathogen – BACTERIA**

	Pathogen	Test result			Laboratory / University	Report Nº
		Medium	Exposure	Effectiveness %		
Hospital Maharakham: Child ICU (Thailand)	BACTERIA					
	Pseudomonas S	Air	4h	85.0 - 100	Green & Healthy Auditors PTY LTD	191009
	Micrococos					
	Escherichia coli					
	Staphylococci					
Aspergillus spp						
Hospital Maharakham: ICU (Thailand)	Pseudomonas S	Air Surface	4h	56.0 - 80.0 70.0 - 100	Green & Healthy Auditors PTY LTD	191120
	Micrococos					
	Staphylococci					
	Aspergillus spp					
Hospital Guangdong (China)	Total of bacteria	Air	24h	68.8 – 78.1	Guang Zhou Institute of Microbiology	KJ20192715
Medical centre Sahmyoook: UCI (korea)	Staphylococcus aureus	Air	2h	53.9	KSD (Korea Securities Depository)	0223190717
	Streptococcus pneumoniae			56.0		
	Escherichia coli			38.7		

## DEPARTMENT OF PATHOLOGY



LILAVATI HOSPITAL  
AND RESEARCH CENTRE

Microbiology section/Department of Pathology  
Lilavati hospital and research center, Mumbai  
Date: 12 September, 2020

## TO WHOMSOEVER IT MAY CONCERN

Subject: Report of the WELLISAIR Air disinfection equipment testing results done at Lilavati Hospital –Pathology Department

## Methods:

1. Swab method
2. Settle plate Method

Areas covered: Microbiology covid testing area

## Procedure:

Pre-Disinfection swabs (Before installing equipment) were taken on 7<sup>th</sup> September, 2020. Then the disinfection equipment was installed in the testing area. After leaving it for 30 minutes swabs were collected from different areas of the section. Swabs were collected each day afternoon from 7<sup>th</sup> to 11<sup>th</sup> September. In addition, Daily settle plates were tested from 7<sup>th</sup> to 11<sup>th</sup> September

## Results-

1. On 7<sup>th</sup> September: Before air disinfection equipment was installed, swab results showed a mild organism growth which was a mixed growth of organisms. However, post installation of equipment, the swabs and settle plates yielded no organism growth.
2. Also, the equipment showed blue signals at most of times during the observation period which indicated sterility of the environment

Name of Equipment : Wellisair Air disinfection		Date of reporting: 12 <sup>th</sup> September					
Date of Testing: 7 <sup>th</sup> to 11 <sup>th</sup> September		Incubator used: Aerobic Incubator					
Sampling method used: Swab & Settle plate		Incubation Temperature: 37 <sup>o</sup> C					
Media Used: Blood agar		Area: Microbiology section of pathology					
Incubation Period: 4 days							
Sampling Date	Pre disinfection/ Post disinfection/ Settle plate	No. of CFU/plate					Average CFU/plates
		On same day	After 24 hours	After 48 hours	After 72 hours	After 96 hours	
7 <sup>th</sup> September	Pre disinfection	20	15	10	10	5	12
	Post disinfection	0	0	0	0	0	0
	Settle plate	0	0	0	0	0	0
8 <sup>th</sup> September	Post disinfection	0	0	0	0	0	
	Settle plate	0	0	0	0	0	
9 <sup>th</sup> September	Post disinfection	0	0	0	0		
	Settle plate	0	0	0	0		
10 <sup>th</sup> September	Post disinfection	0	0	0			
	Settle plate	0	0	0			
Observed by : Dr Rahul Kamble							

Conclusion: "WELLISAIR AIR DISINFECTION EQUIPMENT" has passed the lab testing results in Microbiology department and hence been found satisfactory for use in disinfection of environmental areas

Thanking you

Dr Nitin Chavan  
Consultant Pathologist and Lab Director  
Lilavati Hospital & Research centre, Mumbai

Dr Rahul Kamble  
Consultant Microbiologist & Infection Control Officer  
Lilavati Hospital & Research centre, Mumbai




More than Health Care, Human Care


# WELLISAIR AGAINST BACTERIA: TESTS

Much of the pathogenic microorganisms are part of the family of bacteria. The physical characteristics of bacteria are divided into two groups:

**Bacterium Gram "+":** It is characterized by having two layers on the wall of the bacterium: The external cell wall crossed by protein chains and lipoteichoic acids (among others) and the cytoplasmic membrane (internal wall).



**Bacterium Gram "-":** These bacteria have three layers in the cell wall: The outer membrane composed of phospholipids and lipopolysaccharides, the intermediate layer that is similar to the outer one of the Gram "+" but much thinner, and the inner cytoplasmic membrane the same as the inner membrane of bacteria Gram "+".




The Advanced Oxidation Process (AOP) based on hydroxyl radicals can attack both types of bacteria. This is because both have chains of acids and proteins that are vulnerable to oxidation.


The oxidative reactions that occur cross the bacteria's membranes, modifying their morphological structure (eliminating the membranes), until reaching their genetic information, so that these irreversible damages kill and make the bacteria disappear.

To determine the effectiveness of Wellisair technology against bacteria, various tests were carried out on air and surfaces, such as:

Bacillus Subtilis:




Graph 1. Decrease in the concentration of Bacillus Subtilis plaque forming units per millilitre (PFU / ml) in air.




Graph 2. Decrease in the concentration of Bacillus Subtilis plaque forming units per millilitre (PFU / ml) on the surface 60 minutes after the start of the test.

Staphylococcus Aureus:




Graph 3. Decrease in the concentration of Staphylococcus Aureus plaque forming units per millilitre (PFU / ml) in air.



Graph 4. Decrease in the concentration of Staphylococcus Aureus plaque forming units per millilitre (PFU / ml) on the surface 60 minutes after the start of the test.

Salmonella:



Graph 5. Decrease in the concentration of Salmonella plaque forming units per millilitre (PFU / ml) on the surface 4 hours after the start of the test.


The results of these tests were very satisfactory, because, within a few hours of exposure, 99.9% of the bacteria were eliminated, both in air and on the surface.

# WELLISAIR AGAINST VIRUSES: TESTS

One of the many great ways that Wellisair technology can be used is for virus removal. This process can be divided into two different approaches and viruses: enveloped viruses and non-enveloped viruses. The main difference between them is that enveloped viruses have a protein matrix surrounding the nucleic acid (virus genetic information, RNA). Despite expectations, this matrix is especially sensitive to external agents, such as oxidation.

The Advanced Oxidation Process (POA), which is based on hydroxyl radicals, attacks both types of viruses, but in a different way for each:


**Enveloped virus:** Hydroxyl Radicals react with the fatty acid chains within the envelope (lipid reaction). This reaction causes the membrane to break down, which produces an antioxidant reaction within the virus and accelerates the destruction process (indirect protein oxidation reaction produced by lipid peroxidation) by altering its genetic information.




**Non-enveloped viruses:** Oxidation directly attacks the virus's genetic information (direct protein oxidation reaction produced by lipid peroxidation), making it useless and leaving it on the edge of destruction.

In order to determine the effectiveness of the Wellisair technology, the Laboratory of Virus, Contaminating of Water and Food at the University of Barcelona has carried out a number of tests on different viruses and humidity conditions:

Syncytial RSV:




Graph 1. Reduction in concentration levels of RSV Syncytial plaque forming, units per millilitre (PFU / ml) in dry medium.




Graph 2. Reduction in concentration levels of RSV Syncytial plaque forming, units per millilitre (PFU / ml) in dry medium.

Rotavirus:




Graph 3. Reduction in concentration levels of Rotavirus plaque forming, units per millilitre (PFU / ml) in dry medium.




Graph 4. Percentage reduction of concentration levels of plaque forming, units of Rotavirus in dry medium.

Coxsackievirus:



Graph 5. Reduction in concentration levels of Coxsackievirus plaque forming, units per millilitre (PFU / ml) in dry medium.



Graph 6. Percentage reduction of concentration levels of Coxsackievirus plaque forming, units in dry medium.

The results of these tests demonstrate that Wellisair is effective in fighting against and eliminating viruses, as the oxidation process will behave similarly in all viruses that share a similar morphological structure.

# CERTIFICATES

Patented, tested and certified technology

The innovative patented Wellisair technology, totally harmless to humans, manages to efficiently generate and expand hydroxyl radicals (OH) that, through oxidation, eliminate in a sustainable way 99.9% of viruses, bacteria, allergens, mold, odors and volatiles organic compounds (VOCs) existing in the environment.

The high efficiency in disinfection and purification of the air and surfaces of the Wellisair device has been tested and certified by various studies from prestigious universities both nationally and internationally. It has also been tested in different environments with high risks of virus spread and has been used in sites like WUHAN during the coronavirus crisis.

The following certificates and tests endorse the performance and effectiveness of Wellisair technology.



Universitat de Barcelona  
Laboratory of Viruses Contaminants of Water and Food  
Report on the stability and disinfection of 2019-n-CoV.

The Water and Food Contaminant Virus Laboratory (University of Barcelona) exposed wet and dry viral suspensions to treatment with ozone / d-limonene using the disinfection unit. The result was a reduction of between 99% and 92% of the initial RSV concentration in wet and dry conditions respectively after 2 hours. According to the results in the laboratory, the device can be expected that the efficiency in aerosols and surfaces of the hydroxyl radicals (OH) generated by the tested device, perform a disinfecting action similar to ozone without the inconvenience of toxicity.

**Efficiency against Rotavirus viruses (dry conditions).**

The device was able to reduce 99% of the initial RoV concentration after 2 hours of treatment. Equivalent efficiency is expected receiving equivalent doses of aerosols.

**Efficiency against Coxsackievirus B5 virus.**

The reduction in the number of viruses on dry surfaces ranged from 99.42 to 99.99% after 4 hours.

**Efficiency against RSV virus (Human Respiratory Syncytial Virus) (wet conditions).**

The device was able to reduce 99% of the initial RSV concentration after 2 hours of treatment.

**Efficiency against RSV virus (Human Respiratory Syncytial Virus) (dry conditions).**

The device was able to reduce 92% of the initial RSV concentration after 2 hours of treatment.



Universitat Politècnica de Catalunya  
Laboratory of Sanitary and Environmental Microbiology

**Microbiological test on ozone and peroxide.  
Efficiency against Escherichia coli & Pseudomonas aeruginosa & Staphylococcus aureus bacteria.**

The device was able to reduce more than 99.9% of the initial concentration of Escherichia coli and more than 94% of the initial concentration of Staphylococcus aureus and Pseudomonas aeruginosa microbes after 2 hours of treatment. Both H<sub>2</sub>O<sub>2</sub> work equally in the three bacteria tested.

**Efficiency against Volatile Organic Compounds (VOC) and Particles - CRESCA Test.**

After 30 minutes, without Wellisair the control reaches a harmful air quality and the TVOC and particulate values are above those recommended by international standards. However, with the Wellisair purifier it is possible to maintain good air quality after 30 minutes.



KLC  
Korea Conformity Laboratories  
Efficiency against *Klebsiella pneumoniae* bacteria.

Wellisair air disinfectant was able to reduce 99.9% of the initial airborne concentration of *Klebsiella pneumoniae* microbes after 4 hours of treatment.

Efficiency against bacteria *Staphylococcus Aureus* Resistant to Methicillin (MRSA).

Wellisair air disinfectant was able to reduce 99.9% of the initial airborne concentration of MRSA microbes after 4 hours of treatment.

Efficiency against *Salmonella* & *Klebsiella* & MRSA bacteria

Wellisair air disinfectant was able to reduce 99.9% of the initial microbe concentration of *Salmonella*, *Klebsiella* and MRSA after 4 hours of treatment.

Efficiency against Volatile Organic Compounds (VOCs)

Wellisair air disinfectant achieved a deodorization rate of 80% for ammonia and 99% for formaldehyde after 4 hours of treatment. The deodorization rate was greater than 99.5% for toluene in just 2 hours.

Efficiency against Formaldehydes.

Wellisair air disinfectant achieved a deodorization rate of 19% in the first trial and 41% in the second trial for formaldehyde after treatment.



Kangwon University - Korea

Efficiency against *Bacillus subtilis* & *Escherichia coli* & *Staphylococcus aureus* bacteria (AIR).

The device was able to reduce more than 99.9% of the initial concentration of bacteria in the air of *Escherichia coli* and *Bacillus subtilis* after 20 minutes of treatment and the same reduction of *Staphylococcus aureus* after 60 minutes of treatment.

Efficiency against *Bacillus subtilis* & *Escherichia coli* & *Staphylococcus aureus* bacteria (SURFACE).

The device was able to reduce more than 99% of the initial concentration of surface bacteria after 60 minutes of treatment for *Bacillus subtilis* and *Escherichia Coli* and more than 50% for *Staphylococcus aureus*.



Test Intertek Group plc (New York)

Ozone emissions test

Wellisair meets the criteria required by the standard with a maximum emission of 0.016ppm not higher than the permitted concentration of 0.050ppm. Additionally, it is not necessary to test a second Wellisair per UL867, as the maximum emissions from the first sample have been less than 0.030 ppm, satisfying the exception in section 40.1.1.



- CE Certificates
- CE EMC Certificate
- CE Security Certificate
- FCC EMC Certificate
- Declaration of Conformity RoHS



IJOER  
International Journal of Engineering Research & Science

Evidence of OH· radicals disinfecting indoor air and surfaces in a harmless for humans method



**U.S. Food and Drug Administration (FDA)**



**Laboratory**  
of Viruses Contaminants  
of Water and Food



UNIVERSITAT DE  
BARCELONA

Microbiology, Virology and Biotechnology Section  
Departament de Genètica, Microbiologia i Estadística  
University of Barcelona  
Head of the laboratory: [rgirones@ub.edu](mailto:rgirones@ub.edu)

[http://www.ub.edu/microbiologia\\_virology/en](http://www.ub.edu/microbiologia_virology/en)

## Report on stability and disinfection of 2019-nCoV

The new coronavirus, known as "2019-nCoV", was first detected in Wuhan City (China) and it is presenting a rapid growing number of infected patients. Like other well-known coronaviruses, the primary modes of transmission appear to be direct mucus membrane contact with infectious droplets and through exposure to fomites. This new coronavirus type, has been classified under *Betacoronavirus* genus and, like MERS-CoV and SARS-CoV, has its origins in bats[1].

Several studies proved that closely related coronavirus, SARS-CoV, could be inactivated quite easily with many commonly used disinfectants [2-4]. In particular, SARS-CoV were the most susceptible re-emerging viruses tested against the alcohol-based hand rubs proposed on the WHO guidelines on Hand Hygiene in Health Care [5].

The 2019-nCoV (*Coronaviridae* family) are enveloped viruses about 120 nm in diameter. Like the respiratory syncytial virus (*Pneumoviridae* family) with a 150nm diameter, CoV have surface proteins that mediate the infection of human airway epithelial cells (see figure 1). Those viruses have the ability to survive for many hours on hard surfaces such as tables and crib rails. It typically lives on soft surfaces such as tissues and hands for shorter periods. It is usually transmitted through droplets from the cough or sneeze that contact with eyes, nose, or mouth, or by direct contact with a contaminated surface.

A: *Coronaviridae* (2019-nCoV, MERS and SARS)

B: *Pneumoviridae* (RSV)




Figure 1: Icon representation of the enveloped and spherical viral particles (Viralzone).

The laboratory of viruses contaminants of water and food, tested for WELLIS Co., Ltd, virus stability over time. Wet and dry viral suspensions were exposed to ozone/d-limonene no treatment using the WADU-02, WELLIS disinfection unit. Virus inactivation was calculated against control viral suspensions, not exposed to the disinfection unit and tested in parallel. The disinfection treatment was able to reduce 99% and 92% of the initial concentration of RSV under wet and dry conditions respectively, after 2 hour of treatment. The air disinfection unit WADU-02, WELLIS (Wellis Co.,



Ltd.), significantly reduced the concentration of infectious RSV from wet and dry droplets under laboratory conditions.

It is well known that ozone, at concentrations above 100ppm and high humidity rates, is an effective disinfection treatment, and specially for RNA-viruses with or without envelope [6,7]. However, high ozone concentrations may be harmful to coexist in habitable urban environments. Reactive oxygen species (ROS) including hydroxyl radicals ([OH]), hydrogen peroxide ([H<sub>2</sub>O<sub>2</sub>]) and ozone ([O<sub>3</sub>]) have been reported to enhance disinfection efficiencies of several microorganisms [8,9]. In accordance with the results obtained in our laboratories, it can be expected that the efficiency in aerosols and surfaces that hydroxyl radicals generated by the tested WELLIS WADU-02 device, perform an ozone-like disinfectant action, but without the inconvenience of toxicity.

#### References:

1. **Novel Coronavirus 2019 Situation Summary, Wuhan, China | CDC.** [date unknown],
2. Rabenau HF, Kampf G, Cinatl J, Doerr HW: **Efficacy of various disinfectants against SARS coronavirus.** *J Hosp Infect* 2005, **61**:107–11.
3. Siddharta A, Pfaender S, Vielle NJ, Dijkman R, Friesland M, Becker B, Yang J, Engelmann M, Todt D, Windisch MP, et al.: **Virucidal Activity of World Health Organization–Recommended Formulations Against Enveloped Viruses, Including Zika, Ebola, and Emerging Coronaviruses.** *J Infect Dis* 2017, **215**:902–906.
4. Lai MYY, Cheng PKC, Lim WWL: **Survival of Severe Acute Respiratory Syndrome Coronavirus.** *Clin Infect Dis* 2005, **41**:e67–e71.
5. World Health Organization: **WHO guidelines on hand hygiene in health care: first global patient safety challenge clean care is safer care.** *World Heal Organ* 2009, doi:10.1086/600379.
6. SATO H, WANANABE Y, MIYATA H: **Virucidal Effect of Ozone Treatment of Laboratory Animal Viruses.** *Exp Anim* 1990, **39**:223–229.
7. Breidablik HJ, Lysebo DE, Johannessen L, Skare Å, Andersen JR, Kleiven OT: **Ozonized water as an alternative to alcohol-based hand disinfection.** *J Hosp Infect* 2019, **102**:419–424.
8. Jeong J, Kim JY, Yoon J: **The Role of Reactive Oxygen Species in the Electrochemical Inactivation of Microorganisms.** *Environ Sci Technol* 2006, **40**:6117–6122.
9. Huang X, Qu Y, Cid CA, Finke C, Hoffmann MR, Lim K, Jiang SC: **Electrochemical disinfection of toilet wastewater using wastewater electrolysis cell.** *Water Res* 2016, **92**:164–72.



UNIVERSITAT DE  
BARCELONA

Facultat de Biologia  
Departament de Genètica,  
Microbiologia i Estadística  
Secció de Microbiologia, Virologia  
i Biotecnologia

EZ2911309

04/2019



Yo, MARTA NAVARRO REDONDO, Notario del Ilustre Colegio de CATALUÑA, con residencia en SABADELL.

DOY FE: Que las *mis* fotocopias precedentes que sello y rubrico para su identificación, y la presente que firmo, sello, rubrico y sello, son fiel y exacta reproducción de su original, que tengo a la vista, con el que las otorgo.

Y para que surta todos los efectos legales, expido esta legitimación.

SABADELL, a *8 de junio de 2019*  
Número *54* del libro indicador.



*Marta Navarro Redondo*

## **Wellisair: Eficiencia contra virus Coxackievirus B5**

**Informe de prueba de virus (nº 20180711):** 11 de julio de 2018, Laboratorio de virus contaminantes del agua y de los alimentos de la Universidad de Barcelona.

**Alcance:** medir la eficacia del Wellisair para la desinfección en superficies del Coxsackievirus B5 (CBV5), enterovirus asociado a patologías importantes en humanos como enfermedades de la mano, los pies y la boca e infecciones del sistema nervioso central.

**Procedimiento:** Wellisair fue ubicado en una caja en el interior de la cámara de seguridad a con doce piezas de vidrio contaminadas con 100µl de suspensión viral por muestra a temperatura ambiente. Para los experimentos con virus secos, las piezas de vidrio se secaron durante 1 hora en la vitrina de seguridad antes del experimento de desinfección. Todas las pruebas se realizaron con tres réplicas para cada tiempo y condiciones de tratamiento, y las partículas virales se cuantificaron mediante unidades formadoras de placas (PFU).



### **Resultados:**

<b>Inactivación del CVB5 en suspensiones húmedas</b>			
<b>Tiempo</b>	<b>Sin tratamiento</b>	<b>Tratamiento con Wellisair</b>	
	<b>Virus (PFU/ml)</b>	<b>Virus (PFU/ml)</b>	<b>Reducción (%)</b>
0 min	3,80x10 <sup>8</sup>	3,80x10 <sup>8</sup>	-
30 min	4,10x10 <sup>7</sup>	2,42x10 <sup>7</sup>	40,97
1h	1,33x10 <sup>7</sup>	2,78x10 <sup>6</sup>	79,09
2h	2,17x10 <sup>6</sup>	7,67x10 <sup>1</sup>	99,99
4h	5,27x10 <sup>5</sup>	No se detecta	>99,999

<b>Inactivación del CVB5 en suspensiones secas (primer test)</b>			
<b>Tiempo</b>	<b>Sin tratamiento</b>	<b>Tratamiento con Wellisair</b>	
	<b>Virus (PFU/ml)</b>	<b>Virus (PFU/ml)</b>	<b>Virus (PFU/ml)</b>
0 min	9,87x10 <sup>5</sup>	9,87x10 <sup>5</sup>	-
30 min	1,67x10 <sup>5</sup>	1,00x10 <sup>4</sup>	94,01
1h	1,23x10 <sup>5</sup>	1,07x10 <sup>4</sup>	91,30
2h	4,47x10 <sup>4</sup>	5,47x10 <sup>2</sup>	98,77
4h	1,20x10 <sup>4</sup>	6,90x10 <sup>1</sup>	99,42

<b>Inactivación del CVB5 en suspensiones secas (segundo test)</b>			
<b>Tiempo</b>	<b>Sin tratamiento</b>	<b>Tratamiento con Wellisair</b>	
	<b>Virus (PFU/ml)</b>	<b>Virus (PFU/ml)</b>	<b>Virus (PFU/ml)</b>
0 min	1,53x10 <sup>6</sup>	1,53x10 <sup>6</sup>	-
2h	4,77x10 <sup>5</sup>	2,97x10 <sup>3</sup>	99,39
4h	2,27x10 <sup>5</sup>	1,97x10 <sup>1</sup>	99,99

**Conclusiones:** Wellisair redujo el porcentaje de virus infecciosos CVB5 en superficies húmedas en un 99,99% después de 2 horas de tratamiento y en más del 99,999% después de 4 horas. La reducción del número de virus en superficies secas osciló entre el 99,42 y el 99,99% después de 4 horas.

**Wellisair: Eficiencia contra virus VRS (Virus Sincitial Respiratorio Humano) (condiciones húmedas)**

**Informe de prueba de virus (20191212-3):** 12 de septiembre de 2019, Laboratorio de virus contaminantes del agua y de los alimentos de la Universidad de Barcelona.


**Alcance:** medir la eficacia de Wellisair para la desinfección en superficies contra el virus sincitial respiratorio humano (RSV), principal causa viral de las infecciones agudas del tracto respiratorio inferior: bronquiolitis y neumonía.

**Procedimiento:** Wellisair fue ubicado en una caja de metacrilato junto a cien microlitros en gotas de RSV sobre pequeños trozos de vidrio, humedecidos a temperatura ambiente. En cada tiempo de prueba, los virus se recuperaron mediante medio de cultivo (MEM) y las partículas virales fueron cuantificadas por TCID<sub>50</sub> en células Hep2.



**Resultados:**

Descomposición de la concentración de RSV en condiciones húmedas			
Tiempo	Sin tratamiento	Tratamiento con Wellisair	
	Virus (PFU/ml)	Virus (PFU/ml)	Reducción (%)
0 min	3,58x10 <sup>5</sup>	3,58x10 <sup>5</sup>	-
30 min	2,60x10 <sup>5</sup>	8,08x10 <sup>4</sup>	67%
1h	2,82x10 <sup>5</sup>	5,00x10 <sup>4</sup>	91%
2h	1,77x10 <sup>5</sup>	6,55x10 <sup>3</sup>	99%



**Conclusiones:** el desinfectante de aire Wellisair fue capaz de reducir el 99% de la concentración inicial de RSV después de 2 horas de tratamiento.

Se podría esperar que la eficiencia de Wellisair, recibiendo dosis equivalentes de aerosoles, sea al menos equivalente.

**Wellisair: Eficiencia contra virus VRS (Virus Sincitial Respiratorio Humano)**  
**(condiciones secas)**

**Informe de prueba de virus (nº20191212-4):** 12 de septiembre de 2019, Laboratorio de virus contaminantes del agua y de los alimentos de la Universidad de Barcelona.


**Alcance:** medir la eficacia de Wellisair para la desinfección en superficies contra el virus sincitial respiratorio humano (RSV), principal causa viral de las infecciones agudas del tracto respiratorio inferior: bronquiolitis y neumonía.

**Procedimiento:** Wellisair fue ubicado en una caja de metacrilato junto a cien microlitros en gotas de RSV sobre pequeños trozos de vidrio, secados a temperatura ambiente. Para cada tiempo de prueba, los virus se recuperaron mediante medio de cultivo (MEM) y las partículas virales fueron cuantificadas por TCID<sub>50</sub> en células Hep2.



**Resultados:**

Descomposición de la concentración de RSV en condiciones secas			
Tiempo	Sin tratamiento	Tratamiento con Wellisair	
	Virus (PFU/ml)	Virus (PFU/ml)	Reducción (%)
0 min	5,30x10 <sup>4</sup>	5,30x10 <sup>4</sup>	-
30 min	5,00x10 <sup>4</sup>	1,29x10 <sup>4</sup>	74%
1h	4,30x10 <sup>4</sup>	1,23x10 <sup>4</sup>	87%
2h	4,71x10 <sup>4</sup>	9,05x10 <sup>3</sup>	92%



**Conclusiones:** el desinfectante de aire Wellisair fue capaz de reducir el 92% de la concentración inicial de RSV después de 2 horas de tratamiento.

Se podría esperar que la eficiencia de Wellisair, recibiendo dosis equivalentes de aerosoles, sea al menos equivalente.



**Wellisair: Eficiencia contra Compuestos orgánicos Volátiles (COV) & Partículas**

**Informe de prueba (nº 20190930):** 30 de septiembre de 2019, CRESCA (Universidad Politécnica de Cataluña).

**Alcance:** medir la eficacia de Wellisair para la reducción de los compuestos orgánicos volátiles (COV) y de las partículas atmosféricas en suspensión. Los formaldehidos forman parte de los COV.

**Procedimiento:** la prueba se realizó quemando medio palo de incienso en una cámara con un volumen de 225,72 m<sup>3</sup> (6,6x 6 x 5,7). Las condiciones de calidad del aire se midieron en un primer escenario de control sin ningún tipo de dispositivo purificador de aire y posteriormente con la presencia del Wellisair.

**Resultados:**



**Conclusiones:** después de 30 minutos, sin Wellisair el control llega a una calidad del aire nociva y los valores TVOC & partículas están por encima de lo recomendado por normativas internacionales. Sin embargo, con el purificador Wellisair se consigue mantener una buena calidad del aire a los 30 minutos.

**Wellisair: Eficiencia contra bacteria Klebsiella pneumoniae**

**Informe de prueba de bacterias (nº 027130):** 24 de marzo de 2016, Korea Conformity Laboratories (KCL).

**Alcance:** medir la eficacia de Wellisair para la desinfección contra microbios en el aire de Klebsiella pneumoniae (bacteria causante de infecciones del tracto urinario, neumonía, sepsis, infecciones de tejidos blandos e infecciones quirúrgicas por heridas).

**Procedimiento:**

- Método de prueba: medición de la tasa de reducción de Klebsiella pneumoniae después de inyectar una concentración constante de bacterias dentro de una cámara de ensayos de 8m<sup>3</sup> durante 4 horas.
- Equipo de medición: sistema de toma de aire MAS-100NT.
- Condiciones de prueba: 23°C 50,1% R.H.

**Resultados:**



Antes de tratamiento con Wellisair



Después de tratamiento con Wellisair

<b>Descomposición de la concentración en el aire de Klebsiella pneumoniae</b>			
<b>Tiempo</b>	<b>Antes de tratamiento con Wellisair</b>	<b>Después de tratamiento con Wellisair</b>	
	<b>Microbios (CFU/m<sup>3</sup>)</b>	<b>Microbios (CFU/m<sup>3</sup>)</b>	<b>Reducción (%)</b>
4h	1,2x10 <sup>4</sup>	<10	99,9%

**Conclusiones:** el desinfectante de aire Wellisair fue capaz de reducir el 99,9% de la concentración inicial en el aire de los microbios Klebsiella pneumoniae después de 4 horas de tratamiento.

**Wellisair: Eficiencia contra bacteria Staphylococcus Aureus Resistente a la Meticilina (SARM)**

**Informe de la prueba de bacterias (nº 027131):** 24 de marzo de 2016, Korea Conformity Laboratories (KCL).

**Alcance:** medir la eficacia de Wellisair para la desinfección en el aire contra el Staphylococcus aureus resistente a la meticilina (SARM), bacteria contraída en hospitales a través de la inserción de un tubo de ventilación en el paciente que produce una neumonía nosocomial, una enfermedad que puede ser mortal.

**Procedimiento:**

- Método de prueba: medición de la tasa de reducción de SARM después de inyectar una concentración constante de bacterias dentro de una cámara de ensayos de 8m<sup>3</sup> durante 4 horas.
- Equipo de medición: sistema de toma de aire MAS-100NT.
- Condiciones de prueba: 23°C 50,2% R.H.



Antes de tratamiento con Wellisair



Después de tratamiento con Wellisair

**Resultados:**

Descomposición de la concentración en el aire de SARM			
Tiempo	Antes de tratamiento con Wellisair	Después de tratamiento con Wellisair	
	Microbios (CFU/m <sup>3</sup> )	Microbios (CFU/m <sup>3</sup> )	Reducción (%)
4h	1,0x10 <sup>4</sup>	<10	99,9%

**Conclusiones:** El desinfectante de aire Wellisair fue capaz de reducir el 99,9% de la concentración inicial en el aire de los microbios SARM después de 4 horas de tratamiento.

## Wellisair: Eficiencia contra bacterias Salmonella & Klebsiella & SARM


**Informe de prueba de bacterias (nº 027129):** 24 de marzo de 2016, Korea Conformity Laboratories (KCL).

**Alcance:** medir la eficacia del Wellisair para la desinfección contra **Salmonella Typhimurium** (bacteria causante de diarrea, dolor abdominal, vómitos y náuseas, que puede ser mortal), contra **Klebsiella pneumoniae** (bacteria que causa infecciones del tracto urinario, neumonía, sepsis, infecciones de tejidos blandos e infecciones en heridas quirúrgicas) y contra el **SARM** (bacteria contraída en hospitales a través de la inserción de un tubo de ventilación en el paciente que produce una neumonía nosocomial, enfermedad que puede ser mortal).

### Procedimiento:

- Método de prueba: medición de la tasa de reducción de las diferentes bacterias con una distancia entre el medio inoculado con la cepa y la muestra de 5cm después de 4 horas
- Condiciones de prueba: 37°C 33,1% R.H.

### Resultados:



Elementos de prueba		Resultados del ensayo		
		Inicial (CFU/mL)	Después de 4 h (CFU/mL)	Tasa de reducción (%)
Salmonella Typhimurium	Sin tratamiento	1,6x10 <sup>4</sup>	1,6x10 <sup>4</sup>	-
	Tratamiento con Wellisair	1,6x10 <sup>4</sup>	<10	99,9%
Klebsiella pneumoniae	Sin tratamiento	2,0x10 <sup>4</sup>	2,0x10 <sup>4</sup>	-
	Tratamiento con Wellisair	2,0x10 <sup>4</sup>	<10	99,9%
SARM	Sin tratamiento	1,2x10 <sup>4</sup>	1,2x10 <sup>4</sup>	-
	Tratamiento con Wellisair	1,2x10 <sup>4</sup>	<10	99,9%


**Conclusiones:** el desinfectante de aire Wellisair fue capaz de reducir el 99,9% de la concentración inicial de microbios de Salmonella, Klebsiella y SARM después de 4 horas de tratamiento.

**Wellisair: Emisiones de ozono**


**Informe (n° 103502990CRT-001):** 30 de abril de 2018, Intertek Group plc (Nueva York)

**Alcance:** verificar las emisiones de ozono de Wellisair de acuerdo con la normativa "Electrostatic Air Purifiers, UL867, sección 40, 2016". Para cumplir con los criterios de la norma, la emisión de ozono no puede exceder una concentración de 0,050 ppm.

**Procedimiento:** Wellisair fue ubicado en una cámara de ensayo y se realizaron medidas en cinco ubicaciones con el siguiente equipo: calibrador de ozono Teledyne 703E; monitor de ozono Teledyne 400E; transductor de temperatura Vaisala HMD-70Y y un caudalímetro ST75V. Después de comprobar los dos puntos con concentraciones más altas, las mediciones se repitieron con/sin solución y en modo día/noche durante 8 horas.



**Resultados:**



**Conclusiones:** Wellisair cumple con los criterios requeridos por la norma con una emisión máxima de 0,016ppm no superior a la concentración permitida de 0.050ppm. Además, no es necesario probar un segundo Wellisair según UL867, ya que las emisiones máximas de la primera muestra han sido inferiores a 0,030 ppm, lo que satisface la excepción de la sección 40.1.1.



Product Service

# Attestation of Conformity

No. N8A 096006 0004 Rev. 00

**Holder of Certificate: Wellis Co., Ltd.**W-801, SK V1 Center, 11, Dangsang-ro 41-gil, Yeongdeungpo-gu  
Seoul 07217  
REPUBLIC OF KOREA**Product: Air cleaning appliances  
Air disinfection purifier****Model(s): WADU-02**

<b>Parameters:</b>	Rated voltage:	AC/DC Adapter: 100-240 V~ Air disinfection purifier: DC12 V
	Rated frequency:	AC/DC Adapter: 50/60 Hz
	Rated current:	AC/DC Adapter: 0,5 A
	Protection class:	AC/DC Adapter: Class II Air disinfection purifier: Class III

**Tested according to:** EN 60335-1:2012/A2:2019  
EN 60335-2-85:2003/A11:2012  
EN 62233:2008

This Attestation of Conformity is issued on a voluntary basis according to the Low Voltage Directive 2014/35/EU relating to electrical equipment designed for use within certain voltage limits. It confirms that the listed equipment complies with the principal protection requirements of the directive and is based on the technical specifications applicable at the time of issuance. It refers only to the particular sample submitted for testing and certification. For details see: [www.tuvsud.com/ps-cert](http://www.tuvsud.com/ps-cert)

Test report no.: 077-2286820-000

Date, 2020-07-27

( Brian Cha )

Page 1 of 1

After preparation of the necessary technical documentation as well as the EU declaration of conformity the required CE marking can be affixed on the product. The declaration of conformity is issued under the sole responsibility of the manufacturer. Other relevant EU-directives have to be observed.



Product Service

# Attestation of Conformity

No. N8A 096006 0004 Rev. 00

**Holder of Certificate: Wellis Co., Ltd.**W-801, SK V1 Center, 11, Dangsang-ro 41-gil, Yeongdeungpo-gu  
Seoul 07217  
REPUBLIC OF KOREA**Product: Air cleaning appliances  
Air disinfection purifier****Model(s): WADU-02**

<b>Parameters:</b>	Rated voltage:	AC/DC Adapter: 100-240 V~ Air disinfection purifier: DC12 V
	Rated frequency:	AC/DC Adapter: 50/60 Hz
	Rated current:	AC/DC Adapter: 0,5 A
	Protection class:	AC/DC Adapter: Class II Air disinfection purifier: Class III

**Tested according to:** EN 60335-1:2012/A2:2019  
EN 60335-2-85:2003/A11:2012  
EN 62233:2008

This Attestation of Conformity is issued on a voluntary basis according to the Low Voltage Directive 2014/35/EU relating to electrical equipment designed for use within certain voltage limits. It confirms that the listed equipment complies with the principal protection requirements of the directive and is based on the technical specifications applicable at the time of issuance. It refers only to the particular sample submitted for testing and certification. For details see: [www.tuvsud.com/ps-cert](http://www.tuvsud.com/ps-cert)

Test report no.: 077-2266820-000

Date, 2020-07-27

( Brian Cha )



Product Service

# Attestation of Conformity

No. E8A 096006 0002 Rev. 00

**Holder of Certificate:** **Wellis Co., Ltd.**  
W-801, SK V1 Center, 11, Dangsan-ro 41-gil, Yeongdeungpo-gu,  
Seoul 07217  
REPUBLIC OF KOREA

**Name of Object:** **Air cleaners**  
**(Air Disinfection Purifier)**

**Model(s):** **WADU-02**

**Description of Object:** Rated input voltage of adapter: 110-240 V a.c.  
Rated frequency: 50 / 60 Hz  
Rated output voltage of adapter: 12 V d.c.

**Tested according to:** EN 55014-1:2006/A2:2011  
EN 55014-1:2017  
EN 55014-2:2015  
EN 61000-3-2:2014  
EN 61000-3-3:2013

This Attestation of Conformity is issued on a voluntary basis according to the Directive 2014/30/EU relating to electromagnetic compatibility. It confirms that the listed apparatus complies with all essential requirements of the directive and is based on the technical specifications applicable at the time of issuance. It refers only to the particular sample submitted for testing and certification. For details see: [www.tuvsud.com/ps-cert](http://www.tuvsud.com/ps-cert)

**Test report no.:** KR20-YEC0121

**Date,** 2020-06-03

( Byung-Soo Kang )

Page 1 of 1

After preparation of the necessary technical documentation as well as the EU Declaration of conformity the required CE marking can be affixed on the product. That Declaration of conformity is issued under the sole responsibility of the manufacturer. Other relevant EU-directives have to be observed.

# Supplier's Declaration of Conformity (SDoC)

Report No. : KR20-YEF0044  
Type of equipment : Air Disinfection Purifier  
Model Name : WADU-02  
Unique Identifier : Wellis

## Party issuing Supplier's Declaration of Conformity

Company Name : Wellis Co., Ltd.  
Address : #W-801, SK V1 Center, 11, Dangsan-ro 41-gil,  
Yeongdeungpo-gu, Seoul, Republic of Korea  
Telephone : +82-2-6121-8256  
Contact name : Eunji Kim (e-mail : kimeunzi@welis.co.kr)

## Responsible Party – U.S. Contact Information

Company Name : Wellis Co., Ltd.  
Address : #W-801, SK V1 Center, 11, Dangsan-ro 41-gil,  
Yeongdeungpo-gu, Seoul, Republic of Korea  
Telephone : +82-2-6121-8256  
Contact name : Eunji Kim (e-mail : kimeunzi@welis.co.kr)

## FCC Compliance Statement (for products subject to Part 15)

This device complies with Part 15 of the FCC Rules.

Operation is subject to the following two conditions:

- (1) This device may not cause harmful interference, and
- (2) this device must accept any interference received, including interference that may cause undesired operation.



Approved by/Date:

A handwritten signature in black ink, appearing to read 'NB Jim', is written over a horizontal line.

May. 28, 2020

Department Manager

CE

**DOC**  
Declaration Of Conformity

We

**Wellis Co., Ltd.**

**#W-801, SK V1 Center, 11, Dangsang-ro 41-gil,  
Yeongdeungpo-gu, Seoul, Republic of Korea**

DECLARE ON OUR SOLE RESPONSIBILITY THAT THE PRODUCT

**KIND OF EQUIPMENT : Air Disinfection Purifier**

**TYPE-DESIGNATION : WADU-02**

This EC-Declaration of conformity is following the provisions of RoHS DIRECTIVE 2011/65/EU and its amending.

It is confirmed that a sample of the product has been tested and found in conformity with below standard(s):

**RoHS EN 50581: 2012  
IEC 62321 series (EN 62321 series)**

Detailed specification of the tested product shown in the following test report(s):

**RoHS Test Report : ETLRD200626.0051**

  
**Date of issue :** 30. 09. 2020.

**Authorized By :**


**Wellis Co., Ltd.**  
#W-801, SK V1 Center, 11, Dangsang-ro 41-gil,  
Yeongdeungpo-gu, Seoul, Republic of Korea

**Wellisair: Emisiones de ozono**


**Informe (n° 103502990CRT-001):** 30 de abril de 2018, Intertek Group plc (Nueva York)

**Alcance:** verificar las emisiones de ozono de Wellisair de acuerdo con la normativa "Electrostatic Air Purifiers, UL867, sección 40, 2016". Para cumplir con los criterios de la norma, la emisión de ozono no puede exceder una concentración de 0,050 ppm.

**Procedimiento:** Wellisair fue ubicado en una cámara de ensayo y se realizaron medidas en cinco ubicaciones con el siguiente equipo: calibrador de ozono Teledyne 703E; monitor de ozono Teledyne 400E; transductor de temperatura Vaisala HMD-70Y y un caudalímetro ST75V. Después de comprobar los dos puntos con concentraciones más altas, las mediciones se repitieron con/sin solución y en modo día/noche durante 8 horas.



**Resultados:**



**Conclusiones:** Wellisair cumple con los criterios requeridos por la norma con una emisión máxima de 0,016ppm no superior a la concentración permitida de 0.050ppm. Además, no es necesario probar un segundo Wellisair según UL867, ya que las emisiones máximas de la primera muestra han sido inferiores a 0,030 ppm, lo que satisface la excepción de la sección 40.1.1.



## CERTIFICATE OF REGISTRATION

*This certifies that:*

**WELLIS CO., LTD.  
W801, SK V1 Center  
11 Dangsang-Ro 41-Gil, Yeungdeunpo-Gu  
Seoul , KR 07217**

is registered with the U.S. Food and Drug Administration for FY 2020 pursuant to Title 21, 807 et seq. of the United States Code of Federal Regulations:

Establishment Owner/Operator Number:  
Device Classification Name:

**10077299  
CLEANER, AIR, MEDICAL  
RECIRCULATING**

Product Code:  
Regulation Number:  
Official Correspondent  
and U.S. Agent:

**FRF  
880.5045  
Registrar Corp  
144 Research Drive, Hampton, Virginia,  
23666, USA  
Telephone: +1-757-224-0177 • Fax: +1-757-  
224-0179**


*Registrar Corp will confirm that such registration remains effective upon request and presentation of this certificate until the end of the year stated above, unless said registration is terminated after issuance of this certificate. Registrar Corp makes no other representations or warranties, nor does this certificate make any representations or warranties to any person or entity other than the named certificate holder, for whose sole benefit it is issued. This certificate does not denote endorsement or approval of the certificate-holder's device or establishment by the U.S. Food and Drug Administration. Registrar Corp assumes no liability to any person or entity in connection with the foregoing.*

*Pursuant to 21 CFR 807.39, "Registration of a device establishment or assignment of a registration number does not in any way denote approval of the establishment or its products. Any representation that creates an impression of official approval because of registration or possession of a registration number is misleading and constitutes misbranding."*

*The U.S. Food and Drug Administration does not issue a certificate of registration, nor does the U.S. Food and Drug Administration recognize a certificate of registration. Registrar Corp is not affiliated with the U.S. Food and Drug Administration.*

**Registrar Corp**

144 Research Drive, Hampton, Virginia, 23666, USA  
Telephone: +1-757-224-0177 • Fax: +1-757-224-0179  
info@registrarcorp.com • www.registrarcorp.com

  
**David Lemarz  
Executive Director**

Registrar Corp

Dated: August 24, 2020



## Safety Declaration for Emissions of Hydroxyl Radicals

Considering the standard workplace exposure limit (WEL) currently set at  $1.4 \text{ mg/m}^3$  (1ppm) for Hydrogen Peroxide, the equivalent in hydroxyl radical level is  $(1.4\text{mg/m}^3) / (34\text{g/mol}) = 0.0000411 \text{ mols/m}^3$ .

Calculating figures,  $0.0000411\text{mols/m}^3 \times 6.23 \times 10^{23} = 2.56 \times 10^{19} \text{ molecules/m}^3$ ; if each molecule of Hydrogen Peroxide generates 2 radicals <sup>(1)</sup>,  $2.56 \times 10^{19} \times 2 = 5.12 \times 10^{19} / \text{m}^3$ , **equivalent at  $5.12 \times 10^{13}/\text{ml}$  Radicals as maximum exposure accepted.**

Then calculation for indoor atmosphere with an operational solid Hydrogen Peroxide registered ® used in Air disinfections Models,  $(0.008\text{mols}) \times (6.023 \times 10^{23}) \times 2 = 9.63 \times 10^{21}$  radicals that could be generated, if we consider 60 days the device in continuous operation, then we obtain forming  $1,6 \times 10^{20}$  Radicals per day as average.

If the indoor operating area is  $300\text{m}^3$ ;  $1.6 \times 10^{20}$  radicals per day /  $300 \text{ m}^3$ ,  $= 5.33 \times 10^{17}$  radicals per day /  $\text{m}^3$ , equivalent to  **$5.33 \times 10^{11}/\text{ml}$  Radicals per day**, far away to the limit according the standard mentioned.

Dr. Josep García Raurich  
Director CRESCA-UPC  
Terrassa July 6, 2016

### References

(1) *The Reaction of Ozone with the Hydroxide Ion: Mechanistic Considerations Based on Thermokinetic and Quantum Chemical Calculations and the Role of HO<sub>2</sub> in Superoxide Dismutation. Page2, 6<sup>th</sup> step.* Gábor Merényi, Johan Lind, Sergej Naumov, and Clemens von Sonntag.

**Lab:** Institut Politècnic del Campus de Terrassa (IPCT)  
Ctra. Nacional 150, km 14,5 Sabadell-Terrassa 08220  
**Direcció i gerència:** CRESCA - ESEIAAT - UPC  
C/Colom, 1 - 08222 - Terrassa (Barcelona)  
Tel. +34 3 739 86 54 / +34 3 739 89 86  
<https://www.cresca.upc.edu>

# Evidence of OH• radicals disinfecting indoor air and surfaces in a harmless for humans method

Martínez Vimbert R<sup>1,2</sup>, Arañó Loyo M<sup>1,2</sup>, Custodio Sánchez D<sup>2</sup>, Garcia Raurich J<sup>3</sup>,  
Monagas Asensio P<sup>3</sup>

<sup>1</sup>Emovere Natura, Biomedical Research Communications

<sup>2</sup>e-SHealth, Consulting health care, Parc Científic Universitari Tecno Campus Mataró

<sup>3</sup>Centre de Recerca en Seguretat i Control Alimentari, de la UPC (CRESCA)

**Abstract**— *The development of human societies around the world has generated a very serious environmental damage that threatens human health and the survival of animals and plants due to a higher incidence of infectious diseases.*

*This awareness led to the designing of an advanced harmless environmental sanitation method for the elimination of pathogenic microorganisms and volatile organic compounds (VOC's) in large air spaces and surfaces. Advanced oxidation processes (AOP) based on hydroxyl radicals (OH•) in sufficient concentrations perform biocidal functions on pathogenic microorganisms and degrade airborne organic compounds to mineral forms of harmless organic compounds.*

*It is a technology recognized as clean and safe and is generally carried out through solar radiation as a process initiator with photocatalyst material. The problem presented in the photocatalysis methods is its low speed, the generation of toxic degradation intermediates, deactivation of the material and the need for UV irradiation.*

*The increased airborne spread of pathogenic microorganisms has raised serious concerns about its threat to environmental security. However, there is no effective method to quickly eliminate these harmful microorganisms in a large air space. Compared to conventional disinfectants, OH• radical-based oxidation processes have excellent advantages.*

**Keywords**— *advanced oxidation process, disinfection, hydroxyls radicals (OH•), infectious diseases, ozone, terpenes, VOC's.*

## I. INTRODUCTION

The development of human societies around the world has generated a very serious environmental damage that threatens human health and the survival of animals and plants due to a higher incidence of infectious diseases.

The atmosphere does not have a native microbiota but is a rapid and global means of dispersal for many types of microorganisms. The history of aerobiology has demonstrated until the last century the fundamental role that respiratory air pollution plays in the development of epidemics such as cholera, influenza or Legionella [1]. This respiratory air pollution together with the ease of mobility of human beings around the world has generated in this new century extremely serious respiratory syndromes for survival.

Between November 2002 and July 2003, severe acute respiratory syndrome (SARS) spread rapidly from China to 37 other countries around the world, causing 775 human deaths with an economic loss of \$ 40 billion [2].

In early 2009, a new strain of H1N1 of porcine origin spread worldwide from Mexico. H1N1 was declared a flu pandemic by the World Health Organization (WHO), causing around 17,000 human deaths in early 2010[3].

In 2012 a new episode of coronavirus emerged, the MERS-CoV (Middle East Respiratory Syndrome coronavirus). The emergence of SARS-CoV in 2002 and MERS-CoV in 2012 has changed the perspective of the Coronaviridae family since the pneumonias they have caused (SARS and MERS) have mortality rates of 10% and 30% respectively, which are elevated compared to the rest of the viruses in the family [4].

In December 2019 (after 17 years) a third new coronavirus, named SARS-CoV2 (Severe acute respiratory syndrome coronavirus 2), emerged in Wuhan Hubei province, China [5]. In February 2020, it was renamed as COVID-19 and declared pandemic by the World Organization of Health (WHO). Therefore, it is very important to develop a fast and efficient method for the elimination of pathogenic microorganisms in large air spaces.

This awareness led to the designing of an advanced harmless environmental sanitation method for the elimination of pathogenic microorganisms and volatile organic compounds (VOC's) in large air spaces and surfaces. Advanced oxidation

processes (AOP) based on hydroxyl radicals ( $\text{OH}^\bullet$ ) in sufficient concentrations perform biocidal functions on pathogenic microorganisms and degrade airborne organic compounds to mineral forms of harmless organic compounds [6].

The results of different studies show that  $\text{OH}^\bullet$  radicals rapidly destroy different microorganisms with a concentration of 0.8 mg/L and a spray density of 21  $\mu\text{L}/\text{m}^2$  in 4 seconds [7]. Vital and essential cellular morphological changes in pathogenic microorganisms are also observed under a microscope when exposed to a fatal dose of  $\text{OH}^\bullet$  radicals.

It is a technology recognized as clean and safe and is generally carried out through solar radiation as a process initiator with photocatalyst material. The problem presented in the photocatalysis methods is its low speed, the generation of toxic degradation intermediates, deactivation of the material and the need for UV irradiation.

The increased airborne spread of pathogenic microorganisms has raised serious concerns about its threat to environmental security. However, there is no effective method to quickly eliminate these harmful microorganisms in a large airspace. Compared to conventional disinfectants,  $\text{OH}^\bullet$  radical-based oxidation processes have excellent advantages.

Currently, chlorine, alkali, and alkali-alcohol-amine are the three main types of chemical disinfectants that are widely used to eliminate microbial contamination, but they have some drawbacks. A chemical disinfectant can only selectively kill one or similar types of pathogenic microorganisms; its processing time is long, in a range of 0.5 - 1 hour due to the low chemical reaction rate and a very high lethal dosage value, which could reach 9% (v / v); the remaining chlorine intermediates imply severe secondary contamination. Finally, its lethal processing is limited to the surface of the objects, making it impossible to apply in large air spaces [8,9].

Compared to previous chemical disinfectants, advanced  $\text{OH}^\bullet$  radical-based oxidation technology has several advantages: 1) Absence of selectivity, they can kill any pathogenic microorganism in low lethal doses due to its strong oxidative character, with an oxidation potential of 2.8 V, slightly less than fluorines (3.03 V). 2) The processing time of  $\text{OH}^\bullet$  radicals is very short, several seconds, because the chemical reaction rate of  $\text{OH}^\bullet$  radicals is greater than  $10^9 \text{ L mol}^{-1}\text{second}^{-1}$ , which is  $10^7$  times greater than other oxidants' such as  $\text{O}_3$ ,  $\text{H}_2\text{O}_2$ ,  $\text{Cl}_2$ , etc. 3) As a green oxidant,  $\text{OH}^\bullet$  radicals decompose into  $\text{H}_2\text{O}$  and  $\text{O}_2$  without any residual oxidants after their biochemical reactions [10,11].

## II. MATERIALS AND METHODS

### 2.1 Formation of $\text{OH}^\bullet$ radicals

Oxygen is an essential molecule for life, but due to its high reactivity it also becomes a toxic element that gives rise to the so-called *oxygen paradox*. Oxygen is basically an oxidizing molecule. The following concentration of pollutants is generally found in "clean" outdoor air (without sources of pollution): carbon dioxide, 320 ppm; ozone, 0.02 ppm, carbon monoxide, 0.12 ppm, nitric oxide, 0.003 ppm, and nitrogen dioxide, 0.001 ppm. However, these values increase significantly in urban air [12].

The  $\text{OH}^\bullet$  radical is the most important natural oxidant in tropospheric chemistry, often called the "detergent" in the atmosphere since it reacts with many pollutants, initiating the process of cleaning them up. It also plays an important role in the elimination of greenhouse gases such as carbon dioxide, methane or ozone. Using Advanced Oxidation Processes (AOP) is attractive, among other reasons, because the contaminant is destroyed, not concentrated or transferred to the environment, a total or almost total mineralization of organic pollutants is achieved. Therefore they can be applied in the destruction of the vast majority of organic compounds, especially in non-biodegradable compounds such as organochlorines, PCBs, PAHs, etc. It is a clean and safe technology and in some processes, solar radiation can be used as the initiator.

The main problem for rapidly eliminating pathogenic microorganisms in large air spaces is how to produce the  $\text{OH}^\bullet$  radicals with high concentration and large production. Currently, the main methods are Fenton catalysis, photocatalysis and ozone as well as their collaborative effects [13-17]. However, these technologies have some serious disadvantages: 1) The  $\text{OH}^\bullet$  radicals rate of production is low and they are obtained at low concentration, so that the whole biochemical reaction time is long, in the range of 15 - 360 min. 2) The above-mentioned technologies are only applied to small scale experiments or applications. 3) A large number of chemical reagents such as  $\text{H}_2\text{O}_2$ ,  $\text{TiO}_2$  or  $\text{Fe}^{2+}$ , are necessary in the process of  $\text{OH}^\bullet$  production, resulting in high cost and a safety problem. 4) In order to increase the  $\text{OH}^\bullet$  radicals production, several kinds of technologies are collaborated together resulting in large-volume accessory equipment such as the bubble tower or rotating packed bed.

In previous studies, the production of a large number of OH<sup>•</sup> radicals has been reported by ionization and dissociation of O<sub>2</sub> in air and H<sub>2</sub>O in the gaseous state, using a physical method of strong electric field discharge. In this way, OH<sup>•</sup> radicals have been successfully used in the treatment of ship's ballast water and red-tide in the ocean [18,19].

Titanium oxide is the current reference as a photocatalyst material given its high activity, relative stability, low cost and low toxicity. However, there are problems to be solved such as the low rate of photocatalysis, generation of toxic degradation intermediates, deactivation of the material and the need for UV irradiation as its band gap is not coupled with sunlight [20].

## 2.2 Development of a new advanced oxidation process for the decontamination of air and surfaces.

Given the described scenario, a challenge for safe and effective technological development is generated in the decontamination of air and surfaces. The technological objective is based on the milestone of achieving a method capable of producing OH<sup>•</sup> radicals in sufficient quantities by means of an innovative system that ensures their efficacy and safety for human beings. The Wadu02® system is a device by which are active oxygen species (ROS) such as hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) or a terpene such as d-limonene, evaporates and reacts with an internal ozone emission below a concentration of 0.050 ppm (0.1 mg / m<sup>3</sup>). This ozone exposure limit, established in the regulations issued by the WHO in the environmental limit values (VLA) of the year 2000 for the general public in exposures of up to 8 hours [21], is taken as an international benchmark of safety in ozone emission to obtain a constant and non-damaging production of OH<sup>•</sup> radicals.

Wadu02®'s ozone emissions were evaluated through testing of household electrostatic air cleaners in an external laboratory [22] under the Electrostatic air cleaners standard, SUN - UL 867 and tested with a Teledyne ozone calibrator and monitor, at temperature and humidity controlled by Vaisala transducer and flow meter. The ozone emission of the Wadu02® device was certified in active mode and night mode in parameters less than 0.020ppm (0.012 - 0.015 ppm without filters and 0.015 - 0.016 ppm with filters respectively). Results were lower than all international standards regarding safety in prolonged exposures to ozone.

The production of OH<sup>•</sup> radicals through the oxidation of H<sub>2</sub>O<sub>2</sub> was evaluated under controlled conditions according to the oxidative functionality of the Wadu02® device and compared to liquid hydrogen peroxides in the purity ranges of 0.25% to 0.75%, aided by the colorimetric reaction performed on a potassium iodide test strip. The results indicate that the average oxidative capacity of H<sub>2</sub>O<sub>2</sub> at 0.5% purity is equivalent to the oxidative capacity offered by the Wadu02® model devices, with a maximum production of 0.9 mg/m<sup>3</sup> (0.64 ppm), which is approximately 64.2% of the current workplace exposure limit (WEL) adjusted to 1.4 mg / m<sup>3</sup> (1ppm)[23,24].

The threshold concentration for the acute irritant effects of hydrogen peroxide gas in the respiratory tract is 10 mg/m<sup>3</sup> (equivalent to 7 ppm) in humans, while the corresponding values for the skin are 20mg/m<sup>3</sup>. Regarding its prolonged exposure, hydrogen peroxide has not been found to cause teratogenic or carcinogenic effects in humans. Mutagenic or chromosomal effects have also not been observed.

It was also verified as an alternative to the high natural reactivity of hydrogen peroxide, the substitution of the cartridge load with aromatic essences extracted from flowers and plants due to the biocidal role that terpenes have for their antiviral and antibacterial properties. The process of advanced oxidation was analyzed under the same conditions of low ozone emission (less than 0.02ppm) with the Wadu02® model, to compare the proven efficacy of hydrogen peroxide.

Limonene is one of the most abundant monoterpenes in nature present in essential oils extracted from the peel of citrus fruits, including the essential oils of orange and tangerine. This monoterpene is susceptible to oxidation to generate compounds with higher added value [25].

Terpenes are hydrocarbons present in essential oils that consist of more than one unit of isoprene with five carbons. Monoterpenes, most terpenes, along with sesquiterpenes and diterpenes, comprise the majority of essential oils. Due to the low molecular weight and high volatility of monoterpenes and sesquiterpenes, the use of essential oils in indoor environments can increase the levels of volatile organic compounds (VOCs) [26].

Terpenes contain one or more C=C double bonds, which interact easily with strong oxidants such as ozone, hydroxyl radicals [27-29] and nitrate radicals. Ozone is a common indoor pollutant, the general levels of which are distributed approximately 20 to 40 ppb [30,31]. The use of office machines, such as copy machines, printers and fax machines, also elevates indoor ozone concentrations [32]. The VOCs emitted through the evaporation of indoor terpene-based products may interact with ozone and generate secondary air pollutants, mostly formaldehyde and suspended particulates [33-37]. Secondary organic aerosols generated by the interaction of terpenes and ozone consist of fine and ultrafine particles [38-40]. Consequently, prior

evaluations of total limonene consumption were performed on the Wadu02® device to obtain controlled and safe evaporation.

The total consumption of d-limonene in the Wadu02® products was determined to be of the order of 0.4 g/24 h. According to the functionality of this device and the average evaporation of the measurements recorded in the laboratory, Wadu02® products emit a cloud containing d-limonene with a concentration of approximately 1.84 ppm, which in a room of 60m<sup>2</sup> (180m<sup>3</sup>) can give rise to a maximum concentration with a value less than 2ppb. This concentration is significantly lower than the Swedish and German OEL levels [41] (occupational exposure limits) which are 27ppm and 10ppm, respectively.

### 2.3 Toxicology

For the evaluation and analysis of the amounts of formaldehyde, which can be generated directly from the reaction of ozone with the structural units of C=C bonds, reports indicating that the proportions of formaldehyde formed by this mechanism during ozone-initiated reactions with terpenes represent only a small percentage of reactions to ozone were evaluated [33,34].

The main mechanism that forms formaldehyde is initiated by the reaction of ozone with the functional group C = C to generate ozonide. Subsequently, the ozonide decomposes into a carbonyl and an energy-rich (bi-radical) Criegee intermediate. Both products participate in various additional oxidation reactions to form highly reactive species such as hydroxyl radicals and stable products. These stable products can be ketones and carboxylic acids if the process has taken place in an oxidizing medium, or aldehydes and ketones if the process has taken place in a reducing medium.

The formation of stable carbonyls with low molecular weights, including formaldehyde, acetaldehyde, acetone and propionaldehyde, were observed during the gas-phase reactions of ozone with terpenes [31,33,36]. Reactive hydroxyl radicals generated from ozone reactions with terpenes played a vital role in forming indoor formaldehyde.

Several studies have indicated that indoor OH• radicals concentrations generated by ozone reactions with unsaturated compounds were higher than those outdoors at midday or night [42-44]. OH• radicals were responsible for 56-70% of indoor formaldehyde in reactions between ozone and 23 VOCs and ozone and terpenes [45]. Therefore, a new security objective is the evaluation of the reactions of OH• radicals by means of terpenes and the possible contribution to obtain high levels of formaldehyde indoors and potential effects on indoor air quality [46].

### 2.4 Security test

Once verified that the total consumption of limonene and hydrogen peroxide does not exceed limits considered teratogenic and carcinogenic in humans and that the emission of ozone is less than what is established in international regulations, the effectiveness in reducing formaldehydes was evaluated. The advanced oxidation of limonene with Wadu02® was assessed using the SPS-KACA002-132: 2016 test method under controlled temperature and humidity conditions (21 ± 1) °C (45 ± 5)% RH with d-limonene in the cartridge and with d- limonene in a gel [47]. The results indicate that the reduction of formaldehyde in ozonolysis reactions with emissions less than 0.020 ppm and with low emission concentrations of d-limonene with an evaporation of 0.4 g / 24 h equivalent to 1.84 ppm is significant and reaches values of 19% with gel and 41% with liquid limonene cartridge.

These results show that despite the high reactivity of d-limonene with ozone for the formation of formaldehyde, the controlled emission of ozone below 0.02ppm and the evaporation of limonene below 2ppb in a space of 60m<sup>2</sup> is a safe and harmless reaction.

To validate this hypothesis, a series of experimental tests were carried out to determine the reduction of particles and air pollutants emitted by the burning of an incense stick during a 2-hour exposure to a Wadu02® air purifier, using loaded cartridges with d-limonene and H<sub>2</sub>O<sub>2</sub> in a 225.72 m<sup>3</sup> (6.6 X 6 X 5.7) volume-controlled chamber [48].

Five air quality measurements were performed under different conditions. The first lecture was determined by the initial air quality in the room, without any exposition to incense or air purifiers. The second lecture was taken after 2 hours, since half of an incense stick was burned. The third lecture was determined after the initial air quality was reached and half of an incense stick was burned, with the presence of hydrogen peroxide cartridge air purifier for two hours. The fourth lecture was taken by the same conditions as the third one but, in this case, with a D-limonene cartridge air purifier. Finally, the fifth lecture was determined by the same conditions as third and fourth measurements but, this time, with the presence of both air purifiers (D-limonene and hydrogen peroxide cartridges).



**FIGURE 1. Air quality values before and after contaminating the room with the combustion of an incense stick. Comparison of the safety and efficacy of Wellis in the elimination of VOCs and formaldehyde with hydrogen peroxide and limonene.**

The study shows that under these conditions, burning an incense stick generates poor air quality an average of 30 minutes after the start of combustion with a tendency to regularize after one hour and to return to the initial conditions after two hours, as the particles emitted dispersed in the air space of the room. However, formaldehyde and VOC's readings are higher than the control reading, reflecting the risk of prolonged exposure.

The results in the presence of Wadu02® purifiers regardless of the cartridge content (d-limonene or H<sub>2</sub>O<sub>2</sub>) maintain the initial air quality from the first half-hour of exposure, significantly reducing the values of particle matters, formaldehyde and VOCs. The efficacy in terms of the reduction of formaldehyde and VOCs, according to the use of d-limonene or H<sub>2</sub>O<sub>2</sub> to carry out the emission of OH• radicals, is not significant although H<sub>2</sub>O<sub>2</sub> presents more efficient values.

This allows us to determine that the operation of the Wadu02® air purifier, based on the emission of ozone in low concentrations (< 20 ppb) and the evaporation of standardized amounts of d-limonene or H<sub>2</sub>O<sub>2</sub> from the cartridge (as a result of the execution of the advanced oxidation process) is safe, harmless and effective in reducing suspended particles, VOCs and formaldehyde.

## 2.5 Application of OH• radicals as a broad spectrum biocide

Free radicals and ions cause irreversible alterations in macromolecules (proteins, membranes and DNA) as a consequence of the movement of electrons, resulting in a morbid effect. Reactive Oxygen and Nitrogen Species (RONS) are the most unstable and reactive, meaning these are the first ones to react with others. Within this group, the OH• radicals are the species with a more ephemeral half-life due to its high reactivity, and therefore the most dangerous [49].

The efficacy of the concentration of OH• radicals in the elimination of pathogenic microorganisms was studied. Under conditions of spray density of 21 μL/cm<sup>2</sup> and at processing times of 4 seconds, a dramatic decrease in surviving cells has been reported for *S. Marcescens* at concentrations just above 0.15mg/L and almost entirely in concentrations of 0.41mg/L. In *B. subtilis*, the levels were practically undetectable at concentrations of 0.5 mg/L, whereas in *bacillus* spores the reduction was significant at levels of 0.3mg/L and practically entirely at maximum concentrations of 0.8mg/L [50].

The biocidal function of OH• radicals is based on the advanced oxidation process, a cellular stress mechanism initiated by the "respiratory explosion" (similar to the mitochondrial) and heightened through a cascade of reactions by the release of reactive oxygen species [51-52], such as the hydrogen peroxide, which can pass through biological membranes, and the hypochlorite ion, which modifies and degrades all biological molecules.

The main effects of these reactive forms occur on membranes, lipids and sulfhydryl bonds of DNA's proteins and nucleotides [53], producing:


- Lipid peroxidation, the resulting peroxides of which initiate a catalytic chain reaction leading to further loss of unsaturated fatty acids and extensive membrane damage.
- Production of cross links between proteins, through the formation of disulfide bonds.
- Mutations in the genetic material of the pathogenic microorganism.

During the cellular oxidation process, unsaturated chains are easily attacked by OH• radicals. The peroxidation of the fatty acids in the membranes generates peroxy radicals (ROO•), decreasing their functionality. These radicals have a lower reactivity than the OH• radicals and, therefore, their half-life is somewhat longer.

The presence of cell damage caused by oxidative stress provokes an antioxidant response in the cell: they try to pass electrons from one species to another until radicals are inactive and stability is restored. On the other hand, these interactions can generate cascades spreading the damage [54].

### 2.5.1 Lipid oxidation.

Biological membranes are made up of unsaturated fatty acid chains and are easily oxidized.  $\text{OH}^\bullet$  radicals attack the double bonds of these structures and leave an unpaired electron in the chain that will bind to an oxygen molecule to re-stabilize, giving rise to a peroxy radical. The presence of peroxide radicals modifies the membrane's functionality irreversibly, since it changes its spatial distribution causing instability [55]




**FIGURE 2:  $\bullet\text{OH}$  radicals attack the bonds of these structures and generate a chain with an unpaired electron, which will react with an oxygen molecule ( $\text{O}_2$ ) to re-stabilize. As a product peroxy radicals appear in the membrane, which act as positive feedback further increasing damage.**

### 2.5.2 Protein oxidation: direct (produced by RONS) or indirect (produced by lipid peroxidation)

Free radicals cause changes in the molecular structure of amino acids by modifying their charge, which can end up breaking the polypeptide chain, fragmenting the protein.

Peroxy radicals give rise to substances with aldehyde groups, highly reactive species that establish covalent bridges between amino acids producing intra and interprotein cross-linking. Finally, proteins lose conformation and form aggregates, leading to the decrease or inhibition of the correct functioning of the protein



**FIGURE 3. The  $\text{OH}^\bullet$  radicals form irreversible covalent bridges between two tyrosines. The product (dityrosine) is not recognized by the kinases of the signaling pathways, meaning the information that the tyrosine had to transmit is lost. In addition, this structure is not degradable and therefore non-functional proteins with intra and intermolecular bridges accumulate.**

### 2.5.3 Morphological changes of microorganisms.


The study by Bai et al. 2012 [50] verified the morphological changes of *B. subtilis* and *Bacillus* spores with treatment with  $\text{OH}^\bullet$  radicals under microscopic observation in *B. Subtilis*, *S. Marcescens* and *Bacillus* spores.

It was clearly observed that *B. subtilis* in the form of intact cane and evenly distributed cytoplasm after treatment with  $\text{OH}^\bullet$  radicals, greatly lost the integrity of the membrane. In the other hand, the *Bacillus* spore cells feature a tough, multi-layered outer coating that makes it impossible to quickly kill *Bacillus* spores with conventional chemical disinfectants such as chlorine, alkali and alkali-alcohol-amine. However, after treatment with  $\text{OH}^\bullet$  radicals, the *bacillus* spores also ruptured and the round-shaped cells disappeared. Consequently, *Bacillus* spores require a higher concentration of  $\text{OH}^\bullet$  radicals scavenging, spray density - dispersion and time. Concentration, dispersion spray density and processing time are the three important parameters for the destructive effect of  $\text{OH}^\bullet$  radicals on microorganisms.

The demonstration that the effect of OH• radicals on a microorganism will be with greater biocidal efficacy according to its more superficial structure (collected in previous studies) determines the need to recognize the morphological characteristics of pathogenic microorganisms from their outer layer to the interior of the specific cell [56,57].

Some microorganisms have been able to reverse this oxidative process through superoxide dismutase (SOD), a family of three metalloenzymes (FeSOD, MnSOD and CuZnSOD) with a high capacity to interact with oxidants, neutralizing them and reducing oxidative damage.

MnSOD is synthesized by *Escherichia coli* [58] after exposure to oxygen and is induced by the presence of superoxide radicals. Both SOD and catalase activity have been detected in the cytosol of microbial cells and in the periplasmic space (located between the plasma membrane and the cell wall) of the bacteria. Likewise, a protective role has been demonstrated against ROS generated in the catalase respiratory burst in *Staphylococcus aureus*. [59-61]



**FIGURE 4: Cellular envelopes according to the type and morphology of viruses and bacteria. Adapted from Pearson education, Inc ©2015 & laboratoryinfo.com**

In recent years, different laboratories, external certifiers and university research centers have developed various studies to check the biocidal efficacy of the Wadu02®, in the presence of pathogenic microorganisms in different spaces.


The results are reflected in Table 1, in which we can show that in the case of gram + and gram - bacteria the reduction reaches an average of 99.9% in the first hour of exposure to the advanced oxidation process, in both air and surfaces.

In the case of viruses, the results are observed depending on the conditions of relative humidity and the morphology of the virus. The efficacy results of Wadu02® in non-enveloped viruses indicate that in humid conditions the efficacy is less than in dry environments, averages of 99% are reached; while in enveloped viruses humidity favors the advanced oxidation process and virus elimination than in dry environments.

#### 2.5.4 Report on stability and disinfection of 2019-nCoV

The 2019-nCoV is a new strain of coronavirus that was first detected in Wuhan City (China) in December 2019. The number of infected patients grown rapidly in recent weeks, becoming a serious public health concern. The transmission of the virus occurs mainly via respiratory droplets produced by an infected person that can land in the mouth or nose of people nearby or possibly be inhaled into the lungs. Coronaviruses are a large family of viruses common in many different species of animals, including camels, cattle, cats and bats. Rarely, animal coronaviruses can infect people, and spread among us such as with MERS, SARS and now with 2019-nCoV [62].


The new coronavirus has been classified as a *Betacoronavirus*, like MERS and SARS, both of which have their origins in bats. Coronaviruses are in the subfamily *Coronavirinae* in the family *Coronaviridae*, in the order *Nidovirales*. They are divided in 4 subgenera *Alphacoronavirus*, *Betacoronavirus*, *Deltacoronavirus* and *Gammacoronavirus*[63].



**FIGURE 5: *Nidovirus* phylogeny. The Nidoviral order consists of three families: *Coronavirinae*, *Roniviridae* and *Arteriviridae* from International Committee on Taxonomy of Viruses, 2012 ©.**

Based on the genetic material, these viruses are included in group IV of the Baltimore classification, as the viral particle contains only a single positive-sense stranded RNA. Therefore, the genetic material itself acts as an RNA messenger since they both are positive-sense. When translated, the RNA polymerase and the different structural proteins that form the capsid are synthesized [64].

Coronavirus' diameter is around 60-200 nm. They present a nucleocapsid with helical symmetry and a lipid sheath, that derives from the membrane of the previously infected host cell and contains glycoproteins and surface antigens. From the lipid sheath the characteristic projections of this genus arise forming a solar corona around it that is visible under a microscope and gives the family its name. Despite what might be expected, having an envelope implies that the virus is sensitive to different factors and external agents such as heat, lipid solvents, non-ionic detergents, formaldehyde, oxidizing agents and UV irradiation [63].



**FIGURE 6: Structural proteins of coronavirus (from ViralZone©)**

As can be seen in Figure 6, the coronavirus capsid consists of the following structural proteins:

The Spike glycoprotein (S) protrudes from the outer envelope of the virus forming the "corona" visible under a microscope. Its function is to stick to the proteins found on the cell's surface and infect them. In some cases, the S protein causes the infected cell to fuse with other adjacent cells, thus favoring the spread of the virus. The Envelope protein (E) is responsible for the formation of new viral particles and their release from the infected cell, being necessary for virus diffusion. The Membrane protein (M) is attached to the inner part of the virus membrane and causes this membrane to bend, determining the spherical shape of the virions. M also interacts with the nucleocapsid formed by the RNA of the virus and the N protein. And finally, the Nucleocapsid protein (N) is phosphorylated and binds to the viral genome during assembly.

They are viruses distributed worldwide due to their genetic diversity, their short incubation periods and the high mutation rate they present. The combination of these factors allows the pathogen to infect not just animals but also humans.

It is well known that ozone, at concentrations above 100ppm and high humidity rates, is an effective disinfection treatment especially for RNA-viruses with or without envelope [65-67]. However, high ozone concentrations may be harmful to coexist in habitable urban environments. Reactive oxygen species (ROS) including  $\text{OH}^\bullet$  radicals, hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) and ozone ( $\text{O}_3$ ) have been reported to enhance disinfection efficiencies of several microorganisms [68,69].

### III. RESULTS

According to the results obtained with Wadu02® in viruses with similar structures to Covid-19 (RSV), we can expect that the efficacy of the device will have an expected elimination result of an average of 99 to 92% depending on the relative humidity conditions.

### IV. CONCLUSIONS

The results show that the use of  $\text{OH}^\bullet$  radicals in the advanced oxidation process produced by the Wadu02® purifier is a new, safe and effective method to quickly eliminate pathogenic microorganisms in large air spaces and surfaces.

The application of  $\text{OH}^\bullet$  radicals in different studies has shown that their use in advanced oxidation processes, standardized as a safety measure carried out by Wadu02®, is safe, innocuous and effective in the control of pathogenic microorganisms and the elimination of suspended particles, formaldehyde and VOCs.

The evidence on the efficacy of  $\text{OH}^\bullet$  radicals as a biocide shows that their use is endorsed for being a strong oxidant, capable of eliminating microorganisms in low concentrations (0.8 mg/L) equivalent to 10 thousandths of the dose of conventional

chemical disinfectants. Its spray density - dispersion is  $22\text{ml}/\text{cm}^2$  representing one thousandth of other disinfectants, its constant high reaction rate  $10^9\text{L}/\text{mol} \cdot \text{sec}$  in the processing of  $\text{OH}^\bullet$  radicals is shorter than 4 seconds, which is one thousandth of chemical disinfectants. Finally, the damage that has been observed to pathogens under a microscope is irreversible.

Basing our homeostatic state on the correct functioning of our internal antioxidant system and the experimental demonstration of the use of  $\text{OH}^\bullet$  radicals effectively in the disinfection of air and surfaces, we can issue a safety statement on the use of Wadu02® technology to achieve safe, effective and harmless advanced oxidation processes in humans in the purification and decontamination processes of air and surfaces.

**TABLE 1**  
**WADU02® BIOCIDES EFFICACY TEST RESULT WITH LIMONENE**

Pathogen	Means of dispersion	Exposure	effectiveness %	Documented testing
<i>Bacillus subtilis</i> (Gram +)	Surface	1 h	99,4	Bacillus, Esch, Staph - KNU
	Air	20 min	99,6	Bacillus, Esch, Staph - KNU
<i>Staphylococcus aureus</i> (Gram +)	Surface	1 h	52,3	Bacillus, Esch, Staph - KNU
	Surface	4 h	99,9	Esch, Pseudo, Staphy- KCL
	Air	1 h	99,9	Bacillus, Esch, Staph - KNU
<i>Resistant Staphylococcus aureus (MRSA)</i> (Resistant Gram +)	Surface	4 h	99,9	Salm, Kleb, MRSA - KCL
	Air	4 h	99,9	MRSA - KCL
<i>Pseudomonas aeruginosa</i> (Gram -)	Surface	4 h	99,9	Esch, Pseudo, Staphy- KCL
<i>Enterobacter species: Salmonella</i> (Gram -)	Surface	4 h	99,9	Salm, Kleb, MRSA - KCL
<i>Enterobacter species: Klebsiella</i> (Gram -)	Surface	4 h	99,9	Salm, Kleb, MRSA - KCL
	Air	4 h	99,9	Klebsiella - KCL
<i>Enterobacter species: Escherichia coli</i> (Gram -)	Surface	1 h	99,9	Bacillus, Esch, Staph - KNU
	Surface	4 h	99,9	Esch, Pseudo, Staphy- KCL
	Air	20 min	99,9	Bacillus, Esch, Staph - KNU
<i>Influenza virus</i> (Enveloped)	Wet	30 min	86	Influenza A - UB
	Dry	30 min	38	Influenza A - UB
<i>VRS -Respiratory Syncytial Virus</i> (Enveloped)	Wet	2 h	99	VRS - UB
	Dry	2 h	92	VRS - UB
<i>Rotavirus</i> (Naked)	Wet	2 h	37	RoV - UB
	Dry	2 h	99	RoV - UB

**REFERENCES**

- [1] De la Rosa, MC; Mosso, MA; Ullán, C. (2002) El aire: hábitat y medio de transmisión de microorganismos. *Observ. Medioambiental*; 5:375-402.
- [2] Smith RD. (2006) Responding to global infectious disease outbreaks: lessons from SARS on the role of risk perception, communication and management. *SocSci Med*;63(12):3113-23.
- [3] Writing Committee of the WHO Consultation on Clinical Aspects of Pandemic (H1N1) 2009 Influenza, Bautista E, Chotpitayasunondh T, Gao Z, Harper SA, Shaw M, Uyeki TM, Zaki SR, Hayden FG, Hui DS, Kettner JD, Kumar A, Lim M, Shindo N, Penn C, Nicholson KG. (2010) Clinical aspects of pandemic 2009 influenza A (H1N1) virus infection. *N Engl J Med*; 362(18):1708-19. Review. Erratum in: *N Engl J Med*. 2010 May 27;362(21):2039.
- [4] Chan PK, Chan MC. (2013) Tracing the SARS-coronavirus. *J Thorac Dis*;5(Suppl.2): S118-21.
- [5] Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B. (2020) Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. *Lancet*;395(10223):497-506.
- [6] Isaksen, I.S.A.; Dalsoren, SB. (2011) Getting a better estimate of an atmospheric radical. *Science* 331: 31-9.
- [7] Mindong Bai, Zhitao Zhang, Yiping Tian & Mindi Bai (2012) Rapidly eliminating pathogenic microorganisms in large air space using spraying OH• radicals, *Journal of the Air & Waste Management Association*, 62:4, 393-7.
- [8] Raber, E., Jin, A., Noonan, K., McGuire, R. and Kirvel, R.D. (2001) Decontamination issues for chemical and biological warfare agents: How clean is clean enough? *Int. J. Environ. Health Res.*, 11: 128-148.
- [9] Raber, E. and McGuire, R. (2002) Oxidative decontamination of chemical and biological warfare agents using L-Gel. *J. Hazard. Mater.*, 93: 339-352.
- [10] Roberto, A. and Vincenzo, C. (1999) Advanced oxidation processes (AOP) for water purification and recovery. *Catal. Today*, 53: 51-9.
- [11] Glaze, W.H., Kang, J.W. and Chapin, D.H. (1987) The chemistry of water treatment processes involving ozone, hydrogen peroxide and ultraviolet radiation. *Ozone Sci. Eng.*, 9: 335-6.
- [12] Elías, A; Rogero, S; Fernández, I; Montes, J; Luengas, T; Gurtubay, L. (2012) Contribución de los COV's a la contaminación del aire interior, *Communication XVI International Congress of Project Engineering*. Valencia, Spain.
- [13] Shreya, S., Boris, D. and Vishal, S. (2007) Development of a new approach for microbial decontamination of water using modified Fenton's reaction. *Environ. Pollut.*, 148: 674-8.
- [14] Veressinina, G.Y. and Trapido, M. (2008) Degradation of salicylic acid by Fenton and modified Fenton treatment. *Chem. Eng. J.*, 143: 1-9.
- [15] Maurizio, A. and Vincenzo, A. (2005) Oxidation of oxalate ion in aqueous suspensions of TiO<sub>2</sub> by photocatalysis and ozonation. *Catal. Today*, 107-108: 612-8.
- [16] Naoyuki, K. and Yukako, M. (2005) Advanced oxidation effect of ozonation combined with electrolysis. *Water Res.*, 39: 4661-72.
- [17] Nick, S. and Satoshi, H. (2010) Microwaves in advanced oxidation processes for environmental applications. *J. Photochem. Photobiol. C Photochem. Rev.*, 11: 114-31.
- [18] Bai, M.D., Bai, X.Y., Zhang, Z.T., Bai, M.D. and Yang, B. (2005) Treatment of red tide in Ocean using non-thermal plasma based advanced oxidation technology. *Plasma Chem. Plasma Process.*, 25: 539-50.
- [19] Bai, X.Y., Zhang, Z.T., Bai, M.D. and Yang, B. (2005) Killing of invasive species of ship's ballast water in 20t/h system using hydroxyl radicals. *Plasma Chem. Plasma Process.*, 25: 41-54.
- [20] Gargori, C; Galindo, R; Llusar, M; Tena, M.A; Monrós, G; Badenes, J.A. (2009) Photocatalytic degradation of Orange II by titania addition to sol-gel glasses. *Sci and Tech*; 50:314-20
- [21] WHO (2000) Guías de la calidad del aire de la OMS relativas al material particulado, el ozono, el dióxido de nitrógeno y el dióxido de azufre. (2005) (<http://www.euro.who.int/Document/E87950>)
- [22] Intertek (2018) Ozone emissions testing of household electrostatic air cleaners report number:103502990CRT-001, GFT-OP-10o.
- [23] Real Decreto 374/2001, de 6 de abril sobre la protección de la salud y seguridad de los trabajadores contra los riesgos relacionados con los agentes químicos durante el trabajo. (BOE núm. 104 de 1 de mayo de 2001.)
- [24] Ley 31/1995, de 8 de noviembre, de Prevención de Riesgos Laborales.
- [25] Castellanos, FE. (2007). Biotransformación de limoneno y aceites esenciales de naranja y mandarina empleando aspergillus niger. Chemical Master degree thesis. Universidad Industrial de Santander.
- [26] Bakkali F, Averbeck S, Averbeck D, Waomar M. (2008) Biological effects of essential oils - A review. *Food Chem Toxicol.* 46:446-75.
- [27] Nøjgaard JK, Bilde M, Stenby C, Nielsen OJ, Wolkoff P. (2006) The effect of nitrogen dioxide on particle formation during ozonolysis of two abundant monoterpenes indoors. *Atmos Environ*; 40:1030-42.
- [28] Weschler CJ. (2006) Ozone's impact on public health: Contributions from indoor exposures to ozone and products of ozone-initiated chemistry. *Environ Health Perspect*; 114:1489-96.
- [29] Wells JR, Forester CD. (2009) Yields of carbonyl products from gas-phase reactions of fragrance compounds with OH radical and ozone. *Environ Sci Technol*; 43:3561-8.
- [30] Sabersky RH, Sinema DA, ShairFH. (1973) Concentrations, decay rates, and removal of ozone and their relation to establishing clean indoor air. *Environ Sci Technol*; 7:347-53.

- [31] Weschler CJ. (2000) Ozone in Indoor Environments: Concentration and Chemistry. *Indoor Air*;10:269-88.
- [32] Grosjean D, Williams EL, Grosjean E, Andino JM, Seinfeld JH. Atmospheric oxidation 1 of biogenic hydrocarbons: reaction of ozone with beta-pinene, D-limonene and trans caryophyllene.(1993) *Environ Sci Technol*;27:2754-8.
- [33] Shu Y, Kwok ESC, Tuazon EC, Atkinson R, Arey J. (1997) Products of the gas-phase reactions of linalool with OH radicals, NO<sub>3</sub> radicals, and O<sub>3</sub>. *Environmental science & technology*; 31:896-904.
- [34] Fan Z, Lioy P, Weschler C, Fiedler N, Kipen H, Zhang J. (2003) Ozone-initiated reactions with mixtures of volatile organic compounds under simulated indoor conditions. *Environ Sci Technol*; 37:1811-21.
- [35] Tamás G, Weschler CJ, Toftum J, Fanger PO. (2006) Influence of ozone-limonene reactions on perceived air quality. *Indoor Air*; 16:168-78.
- [36] Lamorena RB, Lee W. (2008) Influence of ozone concentration and temperature on ultra-fine particle and gaseous volatile organic compound formations generated during the ozone-initiated reactions with emitted terpenes from a car air freshener. *J Hazard Mater*; 158:471-7.
- [37] Huang YT, Chen CC, Chen YK, Chiang CM, Lee CY. (2012) Environmental test chamber elucidation of ozone-initiated secondary pollutant emissions from painted wooden panels in buildings. *Build Environ*; 50:135-40.
- [38] Wainman T, Zhang J, Weschler CJ, Lioy PJ. (2000) Ozone and limonene in indoor air: a source of submicron particle exposure. *Environ Health Perspect*; 108:1139-45.
- [39] Lamorena RB, Jung SG, Bae GN, Lee W. (2007) The formation of ultra-fine particles during ozone-initiated oxidations with terpenes emitted from natural paint. *J Hazard Mater*; 141:245-51.
- [40] Sarwar G, Corsi R. (2007) The effects of ozone/limonene reactions on indoor secondary organic aerosols. *Atmos Environ*; 41:959-73.
- [41] Chemical agent directive 98/24/EC and Framework Directive 89/391/EEC
- [42] Weschler CJ, Shields HC. (1996) Production of the hydroxyl radical in Indoor Air. *Environ Sci Technol*; 30:3250-8.
- [43] Weschler CJ, Shields HC. (1997) Measurements of the hydroxyl radical in a manipulated but realistic indoor environment. *Environ Sci Technol*; 31:3719-22.
- [44] Sarwar G, Corsi R, Kimura Y, Allen D, Weschler CJ. (2002) Hydroxyl radicals in indoor environments. *Atmos Environ*; 36:3973-88.
- [45] Weschler CJ. (2004) Chemical reactions among indoor pollutants: what we've learned in the new millennium; 14:184-94.
- [46] Huang, H. L., Tsai, T. J., Hsu, N. Y., Lee, C. C., Wu, P. C., & Su, H. J. (2012). Effects of essential oils on the formation of formaldehyde and secondary organic aerosols in an aromatherapy environment. *Building and Environment*, 57, 120-125.
- [47] Korea Conformity Laboratories (2017). SPS-KACA002-132:2016 / Wadu02-A-03.
- [48] Universitat Politècnica de Catalunya CRESCA-UPC. Salmeron D, AM; Picón T, J; García R, J. (2019) Test of remove effect particle matters, VOC's and formaldehyde by means of hydrogen peroxide and d-limonene.
- [49] Riley PA. (1994) Free radicals in biology: oxidative stress and the effects of ionizing radiation. *Int J Radiat Biol*;65(1):27-33.
- [50] Bai M, Zhang Z, Tian Y, Bai M. (2012) Rapidly eliminating pathogenic microorganisms in large air space using spraying \*OH radicals. *J Air Waste Manag Assoc*; 62(4):393-7.
- [51] Shapoval GS, Gromovaia VF. (2003) Mechanism of antioxidant protection of an organism from oxidative stress. *UkrBiokhim Zh*;75(2):5-13.
- [52] Adair TM. (1996) Is peroxide a hazard? *J Am Dent Assoc*;127(4):422.
- [53] Huttunen J. (1994) Free radicals and antioxidants. Advantages of antioxidants in clinical medicine, truth or fiction? *Duodecim*;110(17):1606-8.
- [54] Ede S, Hafner L, Dunlop P, Byrne J, Will G. (2012) Photocatalytic disinfection of bacterial pollutants using suspended and immobilized TiO<sub>2</sub> powders. *Photochem Photobiol*;88(3):728-35.
- [55] Estrela C, Sydney GB, Bammann LL, Felipe Júnior O. (1995) Mechanism of action of calcium and hydroxyl ions of calcium hydroxide on tissue and bacteria. *Braz Dent J*;6(2):85-90.
- [56] O'Brien D, Stevens N, Fitzgerald-Hughes D, Humphreys H. Effect of a novel air disinfection system on airborne micro-organisms in a hospital outpatient clinic (2012) *J Hosp Infect*;80(1):98-9.
- [57] Wong V, Staniforth K, Boswell TC. (2011) Environmental contamination and airborne microbial counts: a role for hydroxyl radical disinfection units? *J Hosp Infect*;78(3):194-9.
- [58] Hao J, Qiu S, Li H, Chen T, Liu H, Li L. (2012) Roles of hydroxyl radicals in electrolyzed oxidizing water (EOW) for the inactivation of *Escherichia coli*. *Int J Food Microbiol*;155(3):99-104.
- [59] Zhang Q, Ma R, Tian Y, Su B, Wang K, Yu S, Zhang J, Fang J. (2016) Sterilization Efficiency of a Novel Electrochemical Disinfectant against *Staphylococcus aureus*. *Environ Sci Technol*;50(6):3184-92.
- [60] O'Brien D, Stevens N, Fitzgerald-Hughes D, Humphreys H. Effect of a novel air disinfection system on airborne microorganisms in a hospital outpatient clinic. *J Hosp Infect*. 2012 Jan;80(1):98-9.
- [61] Wong V, Staniforth K, Boswell TC. (2011) Environmental contamination and airborne microbial counts: a role for hydroxyl radical disinfection units? *J Hosp Infect*;78(3):194-9.
- [62] Novel coronavirus 2019 situation summary, Wuhan, China, CDC (date unknown)
- [63] Lefkowitz, E. J., Dempsey, D. M., Hendrickson, R. C., Orton, R. J., Siddell, S. G., & Smith, D. B. (2018). Virus taxonomy: the database of the International Committee on Taxonomy of Viruses (ICTV). *Nucleic acids research*, 46(D1), D708–D717.
- [64] Fehr AR, Perlman S. (2015) Coronaviruses: an overview of their replication and pathogenesis. *Methods Mol Biol*; 1282:1-23.
- [65] WHO guidelines on hand hygiene in health care: first global patient safety challenge clean care is safer care

- 
- [66] Sato H, Wananabe, Y; Miyata, H. (1990) Virucidal effect of ozone treatment of laboratory animal viruses. *ExpAnim*; 39:223-9.
- [67] Breidablik, HJ; Lysebo, DE; Johannessen, L; Skare, A; Andersen, JR; Kleiven, OT (2019) Ozonized water as an alternative to alcohol based hand disinfection. *J Hosp Infect*, 102:419-24.
- [68] Jeong, J; Kim, JY; Yoon, J. (2006) The role of Reactive oxygen species in the electrochemical inactivation of microorganisms. *Environ SciTechnol* 40:6117-122.
- [69] Huang, X; Qu, Y; Cid, CA; Finke, C; hoffman, MR; Lim, K; Jiang, SC. (2016) Electrochemical disinfection of toilet wastewater electrolysis cell 92:164-72.